

GAUSSIAN ESTIMATES FOR SCHRÖDINGER PERTURBATIONS

KAROL SZCZYPKOWSKI

We propose a new general method of estimating Schrödinger perturbations of transition densities using an auxiliary transition density as a majorant of the perturbation series. We present applications to Gaussian bounds by proving an optimal inequality involving four Gaussian kernels, which we call 4G Theorem. The applications come with precise control of constants in estimates of Schrödinger perturbations of Gaussian-type heat kernels and also allow for specific non-Kato perturbations.

REFERENCES

- [1] K. Bogdan, W. Hansen and T. Jakubowski. *Localization and Schrödinger perturbations of kernels*. Potential Anal. pages 1–16, 2012.
- [2] K. Bogdan, W. Hansen, and T. Jakubowski. *Time-dependent Schrödinger perturbations of transition densities*. Studia Math., 189 (3):235–254, 2008.
- [3] K. Bogdan, T. Jakubowski, and S. Sydor. *Estimates of perturbation series for kernels*. J. Evol. Equ. 12(4):973–984, 2012.
- [4] K. Bogdan, K. Szczypkowski *Gaussian estimates for Schrödinger perturbations*. Submitted. arXiv:1301.4627v1
- [5] Q. S. Zhang. *On a parabolic equation with a singular lower order term. II. The Gaussian bounds*. Indiana Univ. Math. J. 46(3):989–1020, 1997.
- [6] Q. S. Zhang. *A sharp comparison result concerning Schrödinger heat kernels*. Bull. London Math. Soc. 35(4):461–472, 2003.