

# Escaping points in the boundaries of Baker domains

K. Barański    N. Fagella  
X. Jarque    B. Karpińska

U. Warsaw, U. Barcelona, Technical U. of Warsaw

IMPAN, Simons Semester, Warsaw  
September 16th, 2015

## Fatou components in the escaping set

We consider the dynamical system generated by the iterates of a map

$$f : \mathbb{C} \rightarrow \widehat{\mathbb{C}} \text{ meromorphic (transcendental).}$$

## Fatou components in the escaping set

We consider the dynamical system generated by the iterates of a map

$$f : \mathbb{C} \rightarrow \widehat{\mathbb{C}} \text{ meromorphic (transcendental).}$$

The complex plane decomposes into two totally invariant sets.

- The set of normality of the sequence  $\{f^n\}_n$  is called the **The Fatou set (or stable set)**, denoted by  $\mathcal{F}(f)$ . It is open and its connected components are called **Fatou components**.
- **The Julia set (or chaotic set)**,  $\mathcal{J}(f)$ , is the complement of the Fatou set and closure of the set of repelling periodic points. **Prepoles** are dense in  $\mathcal{J}(f)$ .

## Fatou components in the escaping set

Another relevant set (specially in transcendental dynamics) is the **escaping set**:

$$\mathcal{I}(f) = \{z \in \mathbb{C} \mid f^n(z) \rightarrow \infty\}.$$

## Fatou components in the escaping set

Another relevant set (specially in transcendental dynamics) is the **escaping set**:

$$\mathcal{I}(f) = \{z \in \mathbb{C} \mid f^n(z) \rightarrow \infty\}.$$

For special class of maps (e.g. finite type), the escaping set belongs to the Julia set.

## Fatou components in the escaping set

Another relevant set (specially in transcendental dynamics) is the **escaping set**:

$$\mathcal{I}(f) = \{z \in \mathbb{C} \mid f^n(z) \rightarrow \infty\}.$$

For special class of maps (e.g. finite type), the escaping set belongs to the Julia set.

But in general, there may exist Fatou components which belong to  $\mathcal{I}(f)$ .

Those are:

- (Some) **Wandering domains**  $f^m(U) \cap f^n(U) = \emptyset$  for all  $n, m$ .
- (Some) **Baker domains** Periodic components (period  $k$ ) for which  $\{f^{nk}\}_n$  converge locally uniformly to  $\infty$ . All **invariant** ( $k = 1$ ) Baker domains are in  $\mathcal{I}(f)$ .

# Fatou components in the escaping set

## Natural questions

If  $U$  is a Fatou component in  $\mathcal{I}(f)$ , natural questions are:

- Is  $\partial U$  also in  $\mathcal{I}(f)$ ?

**Answer:** Not in general.

Fatou's example

# Fatou components in the escaping set

## Natural questions

If  $U$  is a Fatou component in  $\mathcal{I}(f)$ , natural questions are:

- Is  $\partial U$  also in  $\mathcal{I}(f)$ ?

Answer: Not in general.

Fatou's example

- Can the opposite occur, i.e.  $\partial U \cap \mathcal{I}(f) = \emptyset$ ?
- In general, how large (in terms of measure) is the set  $\partial U \cap \mathcal{I}(f)$ ?



## Rippon and Stallard's results

These questions were first addressed by Rippon and Stallard.

### Theorem ([RS11], [RS14])

Let  $f$  be an entire transcendental function and let  $U$  be

- an escaping wandering domain, or
- a Baker domain satisfying  $|f^{n+1}(z)| > K|f^n(z)|$  for some  $z \in U$ ,  $K > 1$  and all  $n \geq 1$ , or
- a Baker domain on which  $f$  is univalent.

Let  $\omega$  denote the harmonic measure on  $\partial U$ . Then,  $\omega$ -almost every point in  $\partial U$  escapes.

[RS11] P. J. Rippon and G. M. Stallard, *Boundaries of escaping Fatou components*, Proc. Amer. Math. Soc. 139 (2011), no. 8, 28072820.

[RS14] P.J. Rippon and G. Stallard, *Boundaries of univalent Baker domains*. To appear in J. Anal. Math., 2014

## Further questions

- Is this always the case?

## Further questions

- Is this always the case?
- May the opposite occur?

## Further questions

- Is this always the case?
- May the opposite occur?
- If so, when?

## Further questions

- Is this always the case?
- May the opposite occur?
- If so, when?

To answer these questions we need to briefly introduce

- Inner functions and singularities;
- Classification of Baker domains

# Inner functions

- We work in a slightly more general setting. Consider
  - $f$  a **transcendental meromorphic functions**
  - $U$  a **simply connected** invariant Baker domain.

# Inner functions

- We work in a slightly more general setting. Consider
  - $f$  a **transcendental meromorphic functions**
  - $U$  a **simply connected** invariant Baker domain.
- If  $\varphi : \mathbb{D} \rightarrow U$  is a Riemann map, then the induced map

$$g = \varphi^{-1} \circ f \circ \varphi : \mathbb{D} \rightarrow \mathbb{D}$$

is an **inner function** associated to  $f|_U$ .

# Inner functions

- We work in a slightly more general setting. Consider
  - $f$  a **transcendental meromorphic functions**
  - $U$  a **simply connected** invariant Baker domain.
- If  $\varphi : \mathbb{D} \rightarrow U$  is a Riemann map, then the induced map

$$g = \varphi^{-1} \circ f \circ \varphi : \mathbb{D} \rightarrow \mathbb{D}$$

is an **inner function** associated to  $f|_U$ .

- Clearly

$$\deg(g) = \deg(f|_U),$$

and  $g$  has **no fixed points** in  $\mathbb{D}$ .



## Inner functions

- If  $\deg(g)$  is finite,  $g$  is a proper map (a Blaschke product) and extends to the whole Riemann sphere.

## Inner functions

- If  $\deg(g)$  is finite,  $g$  is a proper map (a Blaschke product) and extends to the whole Riemann sphere.
- If  $\deg(g)$  is infinite, there is at least one **singularity** in  $\partial\mathbb{D}$  (i.e.  $g$  does not extend to any neighborhood of the singular point).

## Inner functions

- If  $\deg(g)$  is finite,  $g$  is a proper map (a Blaschke product) and extends to the whole Riemann sphere.
- If  $\deg(g)$  is infinite, there is at least one **singularity** in  $\partial\mathbb{D}$  (i.e.  $g$  does not extend to any neighborhood of the singular point).
- By the Denjoy-Wolff Theorem, there exists  $p \in \partial\mathbb{D}$  such that

$$g^n \rightarrow p$$

locally uniformly on  $\mathbb{D}$ . The point  $p$  is the **Denjoy-Wolff point** of  $g$ .

## Inner functions

- If  $\deg(g)$  is finite,  $g$  is a proper map (a Blaschke product) and extends to the whole Riemann sphere.
- If  $\deg(g)$  is infinite, there is at least one **singularity** in  $\partial\mathbb{D}$  (i.e.  $g$  does not extend to any neighborhood of the singular point).
- By the Denjoy-Wolff Theorem, there exists  $p \in \partial\mathbb{D}$  such that

$$g^n \rightarrow p$$

locally uniformly on  $\mathbb{D}$ . The point  $p$  is the **Denjoy-Wolff point** of  $g$ .


- We say that  $f|_U$  is **regular** if  $p$  is **NOT a singularity of  $g$** .

# Classification of Baker domains


Baker-Pommerenke-Cowen // König // Barański-F-Jarque-Karpińska

There exists an absorbing domain  $W \subset U$ , and the dynamics in  $W$  are conformally conjugate to either


$$\begin{cases} a\omega, & a > 1 & \text{in } \mathbb{H} & \text{hyperbolic type} \\ \omega \pm i & & \text{in } \mathbb{H} & \text{simply parabolic type} \\ \omega + 1, & & \text{in } \mathbb{C} & \text{doubly parabolic type.} \end{cases}$$



hyperbolic



simply parabolic



doubly parabolic

# Classification of Baker domains

F-Henriksen // Barański-F-Jarque-Karpińska

Classifying particular Baker domains is not an easy task. Some geometric characterizations exist in [FH06] in terms of  $U/f$  and in [BFJK14] in terms of the hyperbolic distance between iterates. The following is relevant for our setting.

# Classification of Baker domains

F-Henriksen // Barański-F-Jarque-Karpińska

Classifying particular Baker domains is not an easy task. Some geometric characterizations exist in [FH06] in terms of  $U/f$  and in [BFJK14] in terms of the hyperbolic distance between iterates. The following is relevant for our setting.

Let  $f$  and  $U$  be as above and let  $\rho_U$  denote the hyperbolic distance in  $U$ .

Theorem ([BFJK14])

$$f|_U \text{ is doubly parabolic} \iff \rho_U(f^{n+1}(z), f^n(z)) \xrightarrow[n \rightarrow \infty]{} 0 \text{ for some } z \in U$$

[BFJK14] K. Barański, N.F. X. Jarque and B. Karpińska, *Absorbing sets and Baker domains for holomorphic maps*, J. of the LMS **92** (2015), 144-162.

# Main Results

We are now ready to state the main results.



## Main Results

We are now ready to state the main results.

Let  $f$  be transcendental meromorphic and  $U$  a simply connected invariant Baker domain.

## Main Results

We are now ready to state the main results.

Let  $f$  be transcendental meromorphic and  $U$  a simply connected invariant Baker domain.

We first see that in many cases, Rippon and Stallard's theorem remains true.

### Theorem A

*Suppose*

- $f|_U$  is hyperbolic or simply parabolic (i.e.  $\rho_U(f^{n+1}(z), f^n(z)) \rightarrow 0$ ), and
- $f|_U$  is regular (e.g. if  $\deg(f) < \infty$ ).

*Then,  $\omega$ -almost every point in  $\partial U$  escapes.*

We remark that if  $f|_U$  is univalent, then it is **always hyperbolic or simply parabolic**.

# Main Results

But then, in the remaining case, **the very opposite occurs**.

# Main Results

But then, in the remaining case, **the very opposite occurs**.

## Theorem B

*Suppose*


- $f|_U$  is doubly parabolic (i.e.  $\rho_U(f^{n+1}(z), f^n(z)) \rightarrow 0$ ), and
- $\deg(f) < \infty$ .

*Then,  $\omega$ -almost every point in  $\partial U$  is topologically recurrent and, in particular, it does NOT escape.*

A point  $z \in \mathbb{C}$  is *topologically recurrent* under  $f$  if its orbit visits any neighborhood of  $z$  infinitely often.

## An Example for Theorem B

Consider  $f(z) = z + e^{-z}$  (Newton's map of  $e^{-e^z}$ ).



- $f$  has infinitely many Baker domains, of degree 2, doubly parabolic.
- Hence the set of escaping points in  $\partial U_i$  has harmonic measure 0.
- We conjecture that
  - all escaping points are nonaccessible from  $U_i$ , while
  - accessible periodic points are dense in  $\partial U_i$ .

## Remarks about Theorem B

- The hypothesis of finite degree **CANNOT** be removed.
  - Aaronson'78 and Doering-Mañé'91 give an example of a simply connected Baker domain, of **infinite degree**, of doubly parabolic type for which  $\omega$ —almost every point in the boundary escapes.

## Remarks about Theorem B

- The hypothesis of finite degree **CANNOT** be removed.
  - Aaronson'78 and Doering-Mañé'91 give an example of a simply connected Baker domain, of **infinite degree**, of doubly parabolic type for which  $\omega$ —almost every point in the boundary escapes.
- On the other hand, there exist Baker domains of infinite degree for which Theorem B holds.

## Remarks about Theorem B

- The hypothesis of finite degree **CANNOT** be removed.
  - Aaronson'78 and Doering-Mañé'91 give an example of a simply connected Baker domain, of **infinite degree**, of doubly parabolic type for which  $\omega$ -almost every point in the boundary escapes.
- On the other hand, there exist Baker domains of infinite degree for which Theorem B holds.
  - Roughly speaking, that happens when

$$\rho_U(f^n(z), f^{n+1}(z)) \longrightarrow 0 \quad \text{fast enough,}$$

even though it is always the case that

$$\sum_{n=0}^{\infty} \rho_U(f^n(z), f^{n+1}(z)) = \infty.$$



## A refinement

The precise condition is as follows, with **no assumption on the degree of  $f|_U$** .

### Theorem C

*Let  $f$  be a meromorphic transcendental map and  $U$  a simply connected invariant Baker domain such that*

$$\rho_U(f^n(z), f^{n+1}(z)) \leq \frac{1}{n} + \mathcal{O}\left(\frac{1}{n^r}\right),$$

*as  $n \rightarrow \infty$  for some  $z \in U$  and  $r > 1$ . Then,  $\omega$ -almost all boundary points are topologically recurrent. In particular, non-escaping points have full harmonic measure.*

## A remark about simply connected parabolic basins

- Suppose  $U$  is an invariant simply connected parabolic basin, i.e. such that  $f^n \rightarrow \zeta \in \partial U \cap \mathbb{C}$  locally uniformly on  $U$ , and  $f'(\zeta) = 1$ .

## A remark about simply connected parabolic basins

- Suppose  $U$  is an invariant simply connected parabolic basin, i.e. such that  $f^n \rightarrow \zeta \in \partial U \cap \mathbb{C}$  locally uniformly on  $U$ , and  $f'(\zeta) = 1$ .
- Consider the analogous set to the escaping set

$$\mathcal{I}_P(f) = \{z \in \partial U \mid f^n(z) \xrightarrow[n \rightarrow \infty]{} \zeta\}.$$

## A remark about simply connected parabolic basins

- Suppose  $U$  is an invariant simply connected parabolic basin, i.e. such that  $f^n \rightarrow \zeta \in \partial U \cap \mathbb{C}$  locally uniformly on  $U$ , and  $f'(\zeta) = 1$ .
- Consider the analogous set to the escaping set

$$\mathcal{I}_P(f) = \{z \in \partial U \mid f^n(z) \xrightarrow[n \rightarrow \infty]{} \zeta\}.$$

- Using extended Fatou coordinates, one can see that  $U$  is of doubly parabolic type in the sense of Baker-Pommerenke-Cowen.

## A remark about simply connected parabolic basins

- Suppose  $U$  is an invariant simply connected parabolic basin, i.e. such that  $f^n \rightarrow \zeta \in \partial U \cap \mathbb{C}$  locally uniformly on  $U$ , and  $f'(\zeta) = 1$ .
- Consider the analogous set to the escaping set

$$\mathcal{I}_P(f) = \{z \in \partial U \mid f^n(z) \xrightarrow[n \rightarrow \infty]{} \zeta\}.$$

- Using extended Fatou coordinates, one can see that  $U$  is of doubly parabolic type in the sense of Baker-Pommerenke-Cowen.
- Then, Theorems B and C remain valid in this setting. In fact, for rational maps, Theorem B was proven in Doering-Mañé'91.

[DM91] Claus I. Doering and Ricardo Mañé, *The dynamics of inner functions.*, Rio de Janeiro: Sociedade Brasileira de Matemática, 1991

## A question remains...

Recall the statement of Theorem C.

### Theorem C

*Let  $f$  be a meromorphic transcendental map and  $U$  a simply connected invariant Baker domain such that*

$$\rho_U(f^n(z), f^{n+1}(z)) \leq \frac{1}{n} + \mathcal{O}\left(\frac{1}{n^r}\right),$$

*as  $n \rightarrow \infty$  for some  $z \in U$  and  $r > 1$ . Then,  $\omega$ -almost all boundary points are topologically recurrent. In particular, non-escaping points have full harmonic measure.*

## A question remains...

Recall the statement of Theorem C.

### Theorem C

*Let  $f$  be a meromorphic transcendental map and  $U$  a simply connected invariant Baker domain such that*

$$\rho_U(f^n(z), f^{n+1}(z)) \leq \frac{1}{n} + \mathcal{O}\left(\frac{1}{n^r}\right),$$

*as  $n \rightarrow \infty$  for some  $z \in U$  and  $r > 1$ . Then,  $\omega$ -almost all boundary points are topologically recurrent. In particular, non-escaping points have full harmonic measure.*

**Question:** Are there actually Baker domains satisfying the assumptions??

## A family of examples

**Answer:** Yes, and in fact there is a whole family of examples.

### Proposition D

Let  $f$  be a meromorphic map of the form

$$f(z) = z + a + h(z)$$


where  $a \in \mathbb{C} \setminus \{0\}$  and

$$|h(z)| < \frac{c_0}{(\operatorname{Re}(z/a))^r} \quad \text{for} \quad \operatorname{Re}\left(\frac{z}{a}\right) > c_1, \quad r > 1, c_0, c_1 > 0.$$

Then  $f$  has an invariant Baker domain  $U$  containing a half-plane  $\{z \in \mathbb{C} : \operatorname{Re}(z/a) > c\}$  for some  $c \in \mathbb{R}$ . Moreover, if  $U$  is simply connected (e.g. if  $f$  is entire), then  $f$  on  $U$  satisfies the assumptions of Theorem C and, consequently,  $\omega$ -almost every point is non-escaping.




## Example 1: Fatou's example




- $f(z) = z + 1 + e^{-z}$  has a Baker  $U$  domain which contains  $\{\operatorname{Re}(z) > 1\}$ .
- The degree of  $f|_U$  is infinite.
- $f$  satisfies the hypothesis of Proposition D (for  $a = 1$ ), hence the Baker domain is doubly parabolic and  $\omega$ -almost every point in  $\partial U$  is topologically recurrent.

## Example 2 ([DM91] and [BFJK15])



- $f(z) = z + \tan z$  has two Baker domains  $U_+ = \{\operatorname{Im}(z) > 0\}$  and  $U_- = \{\operatorname{Im}(z) < 0\}$ .
- The degree of  $f|_{U_{\pm}}$  is infinite.
- $f$  satisfies the hypothesis of Proposition D (for  $a = \pm i$ ), hence the Baker domain is doubly parabolic and  $\omega$ -almost every point in  $\partial U$  is topologically recurrent.

### Example 3: $f(z) = z + i + \tan z$



- (Yellow) Baker domain  $U$  of infinite degree 2. Satisfies the hypothesis of Prop. D for  $a = 2i$ .
- (Black) Infinitely many Baker domains of degree 2, doubly parabolic. Satisfy Theorem B.

## Proof of Theorem A (sketch)

Recall that  $\varphi : \mathbb{D} \rightarrow U$  is a Riemann map,  $g : \mathbb{D} \rightarrow \mathbb{D}$  is the inner function and  $p$  the Denjoy-Wolff point.

## Proof of Theorem A (sketch)

Recall that  $\varphi : \mathbb{D} \rightarrow U$  is a Riemann map,  $g : \mathbb{D} \rightarrow \mathbb{D}$  is the inner function and  $p$  the Denjoy-Wolff point.

- 1 Show that (Lebesgue) almost every point in  $\partial\mathbb{D}$  converges to  $p$ .
  - This is obvious in the univalent case but here one must work locally and extend later.

## Proof of Theorem A (sketch)

Recall that  $\varphi : \mathbb{D} \rightarrow U$  is a Riemann map,  $g : \mathbb{D} \rightarrow \mathbb{D}$  is the inner function and  $p$  the Denjoy-Wolff point.

- 1 Show that (Lebesgue) almost every point in  $\partial\mathbb{D}$  converges to  $p$ .
  - This is obvious in the univalent case but here one must work locally and extend later.
- 2 Prove that almost all of these points map under  $\varphi$  to escaping points.
  - Here we must use a Pflüger type estimate on the behaviour of Riemann maps. This (great) idea is from [RS14].

## Tool 1: A dichotomy

### Theorem[DM91]

Let  $g : \mathbb{D} \rightarrow \mathbb{D}$  be an inner function. Then the following hold.

- (a) If  $\sum_{n=1}^{\infty} (1 - |g^n(z)|) < \infty$  for some point  $z \in \mathbb{D}$ , then  $g^n$  converges to a point  $p \in \partial\mathbb{D}$  almost everywhere on  $\partial\mathbb{D}$ .
- (b) Otherwise  $g$  is recurrent on  $\partial\mathbb{D}$  with respect to the Lebesgue measure.

## Tool 2

This is a quantitative estimate of the principle that “sets that are difficult to reach have very small harmonic measure”.

### Theorem [Pomm92]

Let  $\Phi : \mathbb{D} \rightarrow \mathbb{C}$  be a conformal map, let  $V \subset \Phi(\mathbb{D})$  be a non-empty open set and let  $E$  be a Borel subset of  $\partial\mathbb{D}$ . Suppose that there exist  $\alpha \in (0, 1]$  and  $\beta > 0$  such that:

- (a)  $\text{dist}(\Phi(0), V) \geq \alpha|\Phi'(0)|$ ,
- (b)  $\ell(\Phi(\gamma) \cap V) \geq \beta$  for every curve  $\gamma \subset \mathbb{D}$  connecting 0 to  $E$ .

Then,

$$\lambda(E) < \frac{15}{\sqrt{\alpha}} e^{-\frac{\pi\beta^2}{\text{area}V}}.$$



## Proof of Theorem B (sketch)

- 1 Since  $f|_U$  has finite degree, the inner function  $g$  is a Blaschke product and it extends to the whole sphere.

## Proof of Theorem B (sketch)

- 1 Since  $f|_U$  has finite degree, the inner function  $g$  is a Blaschke product and it extends to the whole sphere.
- 2  $f|_U$  is doubly parabolic, hence  $\mathcal{J}(g) = \partial\mathbb{D}$  [Bargman'08].

## Proof of Theorem B (sketch)

- 1 Since  $f|_U$  has finite degree, the inner function  $g$  is a Blaschke product and it extends to the whole sphere.
- 2  $f|_U$  is doubly parabolic, hence  $\mathcal{J}(g) = \partial\mathbb{D}$  [Bargman'08].
- 3 By results in [Doering-Mañé'91], the series  $\sum_{n=1}^{\infty} (1 - |g^n(z)|)$  is divergent and hence the map  $g$  is recurrent.

## Proof of Theorem B (sketch)


- 1 Since  $f|_U$  has finite degree, the inner function  $g$  is a Blaschke product and it extends to the whole sphere.
- 2  $f|_U$  is doubly parabolic, hence  $\mathcal{J}(g) = \partial\mathbb{D}$  [Bargman'08].
- 3 By results in [Doering-Mañé'91], the series  $\sum_{n=1}^{\infty}(1 - |g^n(z)|)$  is divergent and hence the map  $g$  is recurrent.
- 4  $g$  recurrent implies that  $\omega$ -almost every point in  $\partial U$  is topologically recurrent.

q.e.d.

Thank you for your attention!

# Fatou's Example

Example:  $z \mapsto z + 1 + \exp(-z)$ .



$z = 2k\pi i, k \in \mathbb{Z}$   
are repelling fixed points in  
 $\partial U$  and hence nonescaping.

[Back to questions](#)