

Enhanced Quantization: Solving the Insoluble

John R. Klauder
University of Florida
Gainesville, FL

Enhanced Quantization: The Right Way to Quantize Everything



John R. Klauder
University of Florida
Gainesville, FL




Something Unusual

L. Landau, E.M. Lifshitz, *Quantum mechanics: Non-relativistic theory*, 3rd ed., Pergamon Press, 1977, page 3.

"Thus quantum mechanics occupies a very unusual place among physical theories: it contains classical mechanics as a limiting case, yet at the same time it requires this limiting case for its own formulation."

L & L rule



Classical & Quantum





Planck's constant $h = 6.62606957 \times 10^{-38}$ Joule · sec

$$\hbar = h / 2\pi$$

Class. & Quant. Possibilities




c = classical




q = quantum


Class. & Quant. Possibilities




L & L



L & L



L & L



~~L & L~~

c = classical

q = quantum

“ Triviality of φ_n^4 ”

- Covariant scalar models (imaginary time)

$$I = \int \left(\frac{1}{2} \{ [\nabla \varphi(x)]^2 + m^2 \varphi(x)^2 \} + \lambda \varphi(x)^4 \right) d^n x ; \quad n \geq 5$$

— · — · — · — · — · — · — · — · — · — · — · — · — · — · — · — · —

``It is known that self-interacting scalar fields with a quartic non-linearity do not exist in dimension five or more. (The proofs apply to field theories with a single, scalar field.)”

A. Jaffe and E. Witten (Nov. 2005)



<http://www.claymath.org/sites/default/files/yangmills.pdf>

~~L & L rule~~

that the linear operator H introduced in the preceding section is the energy of the system in quantum mechanics.

In classical mechanics a dynamical system is defined mathematically when the Hamiltonian is given, i.e. when the energy is given in terms of a set of canonical coordinates and momenta, as this is sufficient to fix the equations of motion. In quantum mechanics a dynamical system is defined mathematically when the energy is given in terms of dynamical variables whose commutation relations are known, as this is then sufficient to fix the equations of motion, in both Schrödinger's and Heisenberg's form. We need to have either H expressed in terms of the Schrödinger dynamical variables or H_t expressed in terms of the corresponding Heisenberg dynamical variables, the functional relationship being, of course, the same in both cases. We call the energy expressed in this way the *Hamiltonian* of the dynamical system in quantum mechanics, to keep up the analogy with the classical theory.

A system in quantum mechanics always has a Hamiltonian, whether the system is one that has a classical analogue and is describable in terms of canonical coordinates and momenta or not. However, if the system does have a classical analogue, its connexion with classical mechanics is specially close and one can usually assume that the Hamiltonian is the same function of the canonical coordinates and momenta in the quantum theory as in the classical theory.[†] There would be a difficulty in this, of course, if the classical Hamiltonian involved a product of factors whose quantum analogues do not commute, as one would not know in which order to put these factors in the quantum Hamiltonian, but this does not happen for most of the elementary dynamical systems whose study is important for atomic physics. In consequence we are able also largely to use the same language for describing dynamical systems in the quantum theory as in the classical theory (e.g. to talk about particles with given masses moving through given fields of force), and when given a system in classical mechanics, can usually give a meaning to 'the same' system in quantum mechanics.


Equation (13) holds for v_t any function of the Heisenberg dynamical variables not involving the time explicitly, i.e. for v any constant

[†] This assumption is found in practice to be successful only when applied with the dynamical coordinates and momenta referring to a Cartesian system of axes and not to more general curvilinear coordinates.

DIRAC

The Principles of Quantum Mechanics

Classical \subset Quantum



List of Topics

1 Classical/Quantum connection

“Enhanced Quantization”

Canonical & Affine quantization

Enhanced classical theories



2 A toy model of gravity

3 Ultralocal Quantum Fields

TOPIC 1

- Classical & Quantum formalism
- Canonical coherent states
- Classical \subset Quantum formalism
- Canonical transformations
- Cartesian coordinates
- Affine vs. canonical variables
- Affine quantization as canonical quantization

Action Principle Formulations

Classical action: $A_C = \int [p(t)\dot{q}(t) - H_c(p(t), q(t))] dt$

Variation: $\delta A_C = 0$ yields: $\dot{q} = \partial H_c / \partial p$, $\dot{p} = -\partial H_c / \partial q$

Solution: $p(t)$, $q(t)$ given $p(0), q(0) \in \mathbf{R}^2$

Quantum action: $A_Q = \int \{\langle \psi(t) | [i\hbar\partial/\partial t - \mathcal{H}] | \psi(t) \rangle\} dt$

Variation: $\delta A_Q = 0$ yields $i\hbar\partial|\psi\rangle/\partial t = \mathcal{H}|\psi\rangle$

Solution: $|\psi(t)\rangle$ given $|\psi(0)\rangle \in \mathbf{H}$



VERY DIFFERENT

Restricted Action Principle

Quantum action: $A_Q = \int \{ \langle \psi(t) | [i\hbar\partial/\partial t - \mathcal{H}] | \psi(t) \rangle \} dt$

Possible restrictions: $|\psi(t)\rangle \rightarrow |\psi_E(t)\rangle$ [$\equiv E |\psi(t)\rangle$]

Variation: $\delta A_Q = 0$ yields $[i\hbar\partial|\psi_E(t)\rangle/\partial t = \mathcal{H}_E|\psi_E(t)\rangle]$

$$[\mathcal{H}_E \equiv E \mathcal{H} E]$$

Solution: $|\psi_E(t)\rangle$ given $|\psi_E(0)\rangle \in H$

(1) Nature of $\{|\psi_E(t)\rangle\}$: subspace $\in H$ (**part space**)



(2) Nature of $\{|\psi_E(t)\rangle\}$: subset $\in H$ (**Gaussians**)

Unification of Classical and Quantum (1)

Quantum action: $A_Q = \int \langle \psi(t) | [i\hbar\partial/\partial t - \mathcal{H}] | \psi(t) \rangle dt$

Restricted variation: $|\psi(t)\rangle \rightarrow |\underline{\text{?}}(t)\rangle \in \underline{\mathbf{S}} \subset \mathbf{H}$

sun

Macroscopic variations of Microscopic states:

Basic state:

$$\langle x | \eta \rangle = \eta(x)$$

Translated basic state:

$$\langle x | \eta; q \rangle = \eta(x - q)$$

Translated Fourier state:

$$\langle \kappa | \eta; p \rangle = \tilde{\eta}(\kappa - p)$$

Coherent states:

$$\langle x | p, q \rangle = e^{ip(x-q)/\hbar} \eta(x - q)$$

$$|p, q\rangle \equiv e^{-iqP/\hbar} e^{ipQ/\hbar} |\eta\rangle ; \quad |\eta\rangle = |0\rangle ; \quad (\underline{Q + iP}) |0\rangle = 0$$

14 s.a.

Unification of Classical and Quantum (2)

Quantum action: $A_Q = \int \langle \psi(t) | [i\hbar\partial/\partial t - \mathcal{H}] | \psi(t) \rangle dt$

Restricted variation: $|\psi(t)\rangle \rightarrow |p(t), q(t)\rangle$ **subset**

New action: $A_R = \int \langle p(t), q(t) | [i\hbar\partial/\partial t - \mathcal{H}] | p(t), q(t) \rangle dt$

$$A_R = \int [p(t)\dot{q}(t) - H(p(t), q(t))] dt$$




**NOTE: THIS EQUATION
APPEARS JUST LIKE THE
CLASSICAL ACTION !!!!!**

Unification of Classical and Quantum (2)

Quantum action: $A_Q = \int \langle \psi(t) | [i\hbar\partial/\partial t - \mathcal{H}] | \psi(t) \rangle dt$

Restricted variation: $|\psi(t)\rangle \rightarrow |p(t), q(t)\rangle$ **subset**

New action: $A_R = \int \langle p(t), q(t) | [i\hbar\partial/\partial t - \mathcal{H}] | p(t), q(t) \rangle dt$



$$A_R = \int [p(t)\dot{q}(t) - H(p(t), q(t))] dt$$




CLASSICAL MECHANICS IS QUANTUM
MECHANICS RESTRICTED TO A CERTAIN TWO
DIMENSIONAL SURFACE IN HILBERT SPACE

Canonical Transformations

Restricted quantum action :

$$\begin{aligned} A_R &= \int \langle p, q | [i\hbar\partial/\partial t - \mathcal{H}(P, Q)] | p, q \rangle dt \\ &= \int [p\dot{q} - H(p, q)] dt \end{aligned}$$

Canonical transformations : $p dq = \tilde{p} d\tilde{q} + d\tilde{G}(\tilde{p}, \tilde{q})$

$$\underline{|p, q\rangle} = |p(\tilde{p}, \tilde{q}), q(\tilde{p}, \tilde{q})\rangle \equiv \underline{| \tilde{p}, \tilde{q}\rangle}$$


Restricted quantum action :

$$\begin{aligned} A_R &= \int \langle \tilde{p}, \tilde{q} | [i\hbar\partial/\partial t - \mathcal{H}(P, Q)] | \tilde{p}, \tilde{q} \rangle dt \\ &= \int [\tilde{p}\dot{\tilde{q}} + \dot{\tilde{G}}(\tilde{p}, \tilde{q}) - \tilde{H}(\tilde{p}, \tilde{q})] dt \end{aligned}$$

same quantum



Cartesian Coordinates

Classical/Quantum connection:

$$\begin{aligned} \underline{H(p,q)} &\equiv \langle p,q | \mathcal{H}(P,Q) | p,q \rangle \quad , \quad [(Q + iP) | 0 \rangle = 0] \\ &= \langle 0 | \mathcal{H}(P+p, Q+q) | 0 \rangle = \underline{\mathcal{H}(p,q)} + \mathcal{O}(\hbar; p, q) \end{aligned}$$

Physical meaning: [$\langle 0 | P | 0 \rangle = 0$, $\langle 0 | Q | 0 \rangle = 0$]

$$\boxed{\langle p,q | P | p,q \rangle = p ; \quad \langle p,q | Q | p,q \rangle = q}$$

Fubini-Study metric: [$D_R^2 = \min_{\alpha} \| |\psi\rangle - e^{i\alpha} |\phi\rangle \|^2$]

$$\boxed{2\hbar [\| d | p,q \rangle \|^2 - | \langle p,q | d | p,q \rangle |^2] = dp^2 + dq^2}$$

$$(2/\hbar) [dp^2 \langle \Delta Q^2 \rangle + dp \, dq \langle \{ \Delta Q, \Delta P \} \rangle + dq^2 \langle \Delta P^2 \rangle]$$



Quantum/Classical Summary

Qauntum action :

$$A_Q = \int \langle \psi(t) | [i\hbar \partial / \partial t - \mathcal{H}(P, Q)] | \psi(t) \rangle dt$$

$$\underline{i\hbar \partial | \psi(t) \rangle / \partial t} = \mathcal{H}(P, Q) | \psi(t) \rangle$$

Restricted quantum action :

$$| \psi(t) \rangle \rightarrow | p(t), q(t) \rangle \equiv e^{-iq(t)P/\hbar} e^{ip(t)Q/\hbar} | 0 \rangle$$

$$A_{Q-rest.} = \int \underline{\langle p(t), q(t) |} \underline{[i\hbar \partial / \partial t - \mathcal{H}(P, Q)]} \underline{| p(t), q(t) \rangle} dt$$

$$= \int [p(t)\dot{q}(t) - \underline{H(p(t), q(t))}] dt$$

CT CC
CH \leftarrow QH

$$\dot{q} = \partial H(p, q) / \partial p , \quad \dot{p} = -\partial H(p, q) / \partial q$$

BIG DEAL YIPPEE BIG DEAL HOORAY BIG DEAL



CONVENTIONAL QUANTIZATION RECOVERED



Is There More?

- Are there **other** two-dimensional sheets of normalized Hilbert space vectors that may be used in restricting the quantum action and which lead to an ***enhanced classical canonical formalism?***

Is There More?

- Are there **other** two-dimensional sheets of normalized Hilbert space vectors that may be used in restricting the quantum action and which lead to an **enhanced classical canonical formalism?**

YES !



Affine Variables

Affine variables: $q = q\{q, p\} = \{q, pq\} \equiv \{q, d\}$

$$i\hbar Q = Q[Q, P] = [Q, QP] = [Q, D]; \quad D \equiv (PQ + QP)/2 \quad \text{s.a.}$$

Affine coherent states: $(q > 0 ; \quad Q > 0)$

$$\underline{|p, q\rangle = e^{ipQ/\hbar} e^{-i\ln(q)D/\hbar} |\eta\rangle}; \quad \underline{[(Q-1) + iD/\tilde{\beta}]|\eta\rangle = 0}$$

Overlap function:

$$\langle p', q' | p, q \rangle = \{ \frac{1}{2} [\sqrt{q'/q} + \sqrt{q/q'} + i\sqrt{q'q}(p'-p)/\tilde{\beta}] \}^{-2\tilde{\beta}/\hbar}$$

Resolution of unity:

$$I = \int |p, q\rangle \langle p, q| dp dq / 2\pi\hbar / (1 - 1/2\tilde{\beta}/\hbar)$$

→ **also** $(q < 0, \quad Q < 0)$ **U** $(q > 0, \quad Q > 0)$ ← 23

Affine Quantization (1)

Quantum action :

$$A_Q \equiv \int \langle \psi(t) | [i\hbar\partial/\partial t - \mathcal{H}(D, Q)] | \psi(t) \rangle dt$$

Restricted action : $|\psi(t)\rangle \rightarrow |p(t), q(t)\rangle$ **subset**

$$A_R \equiv \int \langle p(t), q(t) | [i\hbar\partial/\partial t - \mathcal{H}(D, Q)] | p(t), q(t) \rangle dt$$

$$= \int [-q(t) \dot{p}(t) - H(p(t), q(t))] dt$$



Canonical transformation : $|\tilde{p}, \tilde{q}\rangle = |p, q\rangle$

$$A_R \equiv \int \langle \tilde{p}(t), \tilde{q}(t) | [i\hbar\partial/\partial t - \mathcal{H}(D, Q)] | \tilde{p}(t), \tilde{q}(t) \rangle dt$$

$$= \int [-\tilde{q}(t) \dot{\tilde{p}}(t) + \dot{\tilde{G}}'(\tilde{p}(t), \tilde{q}(t)) - \tilde{H}(\tilde{p}(t), \tilde{q}(t))] dt$$

Affine Quantization (2)

Classical/Quantum connection:

$$\begin{aligned}\underline{H'(pq,q)} &\equiv \langle p,q | \mathcal{H}(D,Q) | p,q \rangle , \quad [(Q-1) + iD / \tilde{\beta}] |\eta\rangle = 0 \\ &= \langle \eta | \mathcal{H}(D + pqQ, qQ) | \eta \rangle = \underline{\mathcal{H}(pq,q)} + \mathcal{O}(\hbar; p, q)\end{aligned}$$

Physical meaning: [$\langle \eta | D | \eta \rangle = 0$, $\langle \eta | Q | \eta \rangle = 1$]

$$\langle p,q | D | p,q \rangle = pq ; \quad \langle p,q | Q | p,q \rangle = q$$

Fubini-Study metric:

$$2\hbar [\|d|p,q\rangle\|^2 - |\langle p,q | d | p,q \rangle|^2] = \underline{\tilde{\beta}^{-1}q^2 dp^2 + \tilde{\beta} q^{-2} dq^2}$$

Poincaré half plane: geodesically complete

The Q/C Connection : Summary

- *The classical action arises by a restriction of the quantum action to coherent states*
- Canonical quantization uses P and Q which must be self adjoint
- Affine quantization uses D and Q which are self adjoint when $Q > 0$ (and/or $Q < 0$)
- *Both canonical AND affine quantum versions are consistent with classical, canonical phase space variables p and q*

— · — · — · — · — · — · —

- *Now for some applications!*

TOPIC 2

- A toy model of gravity has singularities

$$A_C = \int_0^T [-q(t) \dot{p}(t) - q(t) p(t)^2] dt , \quad q(t) > 0$$

$$\dot{p}(t) = -p(t)^2$$

$$p(t) = p_0(1 + p_0 t)^{-1} , \quad q(t) = q_0(1 + p_0 t)^2$$

- Canonical quantum corrections
- Affine quantum corrections
- *Affine quantization resolves singularities!*

Toy Model

Classical action.: $A_C = \int [-q\dot{p} - qp^2] dt ; \quad q > 0$

Solution: $p(t) = p_0(1 + p_0 t)^{-1}, \quad q(t) = q_0(1 + p_0 t)^2$

Canonical quant.: $\langle p, q | P Q P | p, q \rangle = qp^2 + qa^2 ; \quad a^2 = \hbar/2$

Solution: $p(t) = a \cot(a(t + \tau)), \quad q(t) = (E_0/a^2) \sin(a(t + \tau))^2$

Affine quant.: $\langle p, q | D Q^{-1} D | p, q \rangle = qp^2 + \hbar^2 C / q \quad \leftarrow$

Solution: $p(t) = \frac{(t + \tau)}{(t + \tau)^2 + K}, \quad \underline{q(t) = M[(t + \tau)^2 + K] > 0}$



$$\hbar^2 C = \langle \eta | D Q^{-1} D | \eta \rangle; \quad K = \hbar^2 C / 4 E_0^2; \quad M = 4 E_0$$

$$\rightarrow e^{iqP/\hbar} Q e^{-iqP/\hbar} = Q + q; \quad e^{i \ln(q) D / \hbar} Q e^{-i \ln(q) D / \hbar} = q Q \quad \leftarrow$$

Enhanced Toy Model : Summary


- A toy model of gravity exhibits singular solutions for all positive energies
- *Enhanced classical theory with affine quantum corrections removes all singularities*
- *Enhanced quantization can eliminate classical singularities (Works for cosmological models)*

TOPIC 3

- Ultralocal scalar field models $[x \in \mathbf{R}^s]$

$$A = \int \left\{ \frac{1}{2} [\dot{\varphi}(t, x)^2 - m_0^2 \varphi(t, x)^2] - g_0 \varphi(t, x)^4 \right\} dt dx$$

- Non-renormalizable quantum theory



$$(p_0^2 + \cancel{\vec{p}}^2 + m_0^2)^{-1}$$

- Also a trivial (=free) theory
- Affine quantization is the key idea
- But first, two important remarks

Free & Pseudofree Theories - 1




Classical action :

$$A_g(x) = A_0(x) + gA_I(x) ; \quad \lim_{g \rightarrow 0} A_g = ?$$

Example 1: $A_g = \int \left\{ \frac{1}{2} [\dot{x}(t)^2 - x(t)^2] - gx(t)^4 \right\} dt$

Example 2: $A_g = \int \left\{ \frac{1}{2} [\dot{x}(t)^2 - x(t)^2] - gx(t)^{-4} \right\} dt$

Moral: $\lim_{g \rightarrow 0} A_g \equiv A'_0$; (1) $A'_0 = A_0$, (2) $A'_0 \neq A_0$



Free & Pseudofree Theories - 2

Free action :

$$A_0 = \int \left\{ \frac{1}{2} [\dot{x}^2 - x^2] \right\} dt$$

Interacting action : $A_g = \int \left\{ \frac{1}{2} [\dot{x}^2 - x^2] - gx^{-4} \right\} dt$

Pseudofree action : $\lim_{g \rightarrow 0} A_g \equiv A'_0 \neq A_0$

Free quantum propagator :

$$\begin{aligned} K_f(x'', T; x', 0) &= N_0 \int e^{-\int \left\{ \frac{1}{2} [\dot{x}^2 + x^2] \right\} dt} Dx \\ &= \sum_{n=0}^{\infty} h_n(x'') h_n(x') e^{-(n+1/2)T} \end{aligned}$$

Pseudofree quantum propagator :

$$\begin{aligned} K_{pf}(x'', T; x', 0) &= \lim_{g \rightarrow 0} N_g \int e^{-\int \left\{ \frac{1}{2} [\dot{x}^2 + x^2] + gx^{-4} \right\} dt} Dx \\ &= \theta(x'' x') \sum_{n=0}^{\infty} h_n(x'') h_n(x') [1 - (-1)^n] e^{-(n+1/2)T} \end{aligned}$$

Ulralocal Scalar Models (1)

Classical action : $[x \in \mathbf{R}^s]$

$$A = \int \{\pi\dot{\phi} - \frac{1}{2}[\pi^2 + m_0^2\phi^2] - g_0\phi^4\} dt dx$$

Free quantum model on a lattice :

$$C(f) = M \int e^{i \sum_k f_k \phi_k a^s - m_0 \sum_k \phi_k^2 a^s} \prod_k d\phi_k \rightarrow e^{-\int f(x)^2 dx / 4m_0}$$

Interacting model on a lattice : **[C.L.T.]**

$$C(f) = M \int e^{i \sum_k f_k \phi_k a^s - \sum_k Y(\phi_k^2, a)} \prod_k d\phi_k \rightarrow e^{-\int f(x)^2 dx / 4\mathbf{m}}$$

Non-renormalizable AND Trivial

Central Limit Theorem

$$C(h) = " \mathcal{V} \int \exp \{ i \int h(x) \varphi(x) dx - \int G(\varphi(x)^2) dx \} \prod_x d\varphi(x) "$$


$$\tilde{C}_K(h) = \tilde{N} \int \exp \left\{ \sum_{k=1}^K [ih_k \varphi_k \Delta - G(\varphi_k^2) \Delta] \right\} \prod_{k=1}^K d\varphi_k$$

$$\begin{aligned} C_K(h) &= N \prod_{k=1}^K \int \exp [ih_k u - G(\Delta^{-2} u^2) \Delta - F(\hbar, u^2, \Delta)] du \\ &= \prod_{k=1}^K [1 - h_k^2 \langle u^2 \rangle / 2! + h_k^4 \langle u^4 \rangle / 4! - h_k^6 \langle u^6 \rangle / 6! + \dots] \end{aligned}$$


Lim $K \rightarrow \infty$ requires $\langle u^2 \rangle = O(\Delta)$, which can be done in TWO basic ways :

1) $\langle u^{2m} \rangle = O(\Delta^m)$, leads to $C(h) = \exp[-A \int dx h(x)^2]$ (C.L.T.)

2) $\langle u^{2m} \rangle = O(\Delta)$, leads to $C(h) = \exp(-\int dx \int \{1 - \cos[uh(x)]\} W(u) du)$



1



2

Ulralocal Scalar Models (2)

Affine quantization : $[\hat{\phi}(x), \hat{\kappa}(y)] = i\delta(x - y)\hat{\phi}(x)$

$$\underline{\pi(x)^2} = \{\pi(x)\phi(x)\}\phi(x)^{-2}\{\phi(x)\pi(x)\} \equiv \kappa(x)\phi(x)^{-2}\kappa(x)$$

$$\rightarrow \hat{\kappa}(x)\hat{\phi}(x)^{-2}\hat{\kappa}(x) = \underline{\hat{\pi}(x)^2 + \frac{3}{4}\hbar^2\delta(0)^2\hat{\phi}(x)^{-2}}$$

using $\hat{\kappa}(x) \equiv \frac{1}{2}[\hat{\pi}(x)\hat{\phi}(x) + \hat{\phi}(x)\hat{\pi}(x)]$

Lattice Hamiltonian : $F \equiv (\frac{1}{2} - ba^s)(\frac{3}{2} - ba^s)a^{-2s}$

$$\mathcal{H} = \frac{1}{2}\sum_k \{-\hbar^2 a^{-2s} \partial^2 / \partial \phi_k^2 + m_0^2 \phi_k^2 + 2g_0 \phi_k^4 + \hbar^2 F \phi_k^{-2} - E_0\} a^s$$

includes a novel counter term !

Ultralocal Scalar Models (3)

Interacting model ground state distribution :

$$C(f) = \int \prod_k (ba^s) \{ e^{if_k \phi_k a^s - Y(\phi_k^2, b, a)} |\phi_k|^{-(1-2ba^s)} d\phi_k \}$$
$$\rightarrow e^{-b \int dx [1 - \cos(f(x)\lambda)]} \exp[-y(\lambda^2, b)] d\lambda / |\lambda|$$


Pseudofree distribution : $(g_0 \rightarrow 0)$

$$C_{pf}(f) = e^{-b \int dx [1 - \cos(f(x)\lambda)]} \exp[-bm\lambda^2] d\lambda / |\lambda|$$

$2m, 0$

Theory space

Free o



Ulralocal Scalar Models (4)

$$A_g = \int \left\{ \frac{1}{2} [\dot{\phi}(t, x)^2 - m_0^2 \phi(t, x)^2] - g_0 \phi(t, x)^4 \right\} dt d^s x$$

Free: $I_p = M \int [\sum'_k \phi_k^2 a^s]^p e^{-m_0 \sum'_k \phi_k^2 a^s} \prod'_k d\phi_k = O(N'^p) \rightarrow \infty$

Hyper - spherical coordinates : $\boxed{\phi_k = \kappa \eta_k ; \quad \kappa^2 = \sum'_k \phi_k^2}$

$$I_p = M \int [\kappa^2 a^s]^p e^{-m_0 \kappa^2 a^s} \kappa^{(N'-1)} d\kappa d\mu(\eta) = O(N'^p) \rightarrow \infty$$

.....

Pseudofree: $\kappa^{(N'-1)} \rightarrow \kappa^{(R-1)} ; \quad R = 2ba^s N' < \infty$

$$J_p = M' \int [\sum'_k \phi_k^2 a^s]^p e^{-m_0 \sum'_k \phi_k^2 a^s} \prod'_k [\phi_k^2]^{-(1-2ba^s)/2} \prod'_k d\phi_k < \infty$$

NO DIVERGENCES

Ultralocal Models : Summary

- Canonical quantization of interacting ultralocal scalar fields is **perturbatively non-renormalizable** and **rigorously trivial**
- Affine quantization of interacting ultralocal scalar fields is **rigorously nontrivial** & **NO** divergences
- Ultralocal scalar models involve **discontinuous perturbations** for which interacting theories are continuously connected to a **pseudofree theory** and **not** to their own **free theory**
- **Hyper-spherical radius variable measure is a simple key to solution** : $\kappa^{N-1} \rightarrow \kappa^{R-1}$; $R < \infty$



Canonical vs. Enhanced

- Canonical quantization requires Cartesian coordinates, but **WHY** is not clear
 - Canonical quantization works well for many problems, but **NOT** for all problems
- · — · — · — · — · — · — · — · — · — · — · — · — · — · —

- Enhanced quantization clarifies coordinate transformations and Cartesian coordinates
- Enhanced quantization can yield canonical results -- **OR** provide proper results when canonical quantization fails

Main Message of Today

**Class./Quan. Coexistence *IS*
Possible *AND IT IS* Beneficial**



GOOD

BETTER



Enhanced Quantization:

Additional Models

Resolved by *EQ*



John R. Klauder
University of Florida
Gainesville, FL



List of Topics

- 1 A toy model of hydrogen
- 2 Enhanced quantization of spin
- 3 Rotationally symmetric models
- 4 Sketch of affine quantum gravity
- 5 AHS model of spikes

Toy Model

Affine quantization: $[Q, P]Q = i\hbar Q = \underline{[Q, D]} , \quad D = \frac{1}{2}(QP + PQ)$

Affine coherent states: $|p, q\rangle = e^{ipQ/\hbar} e^{-i\ln(q)D/\hbar} |\eta\rangle ; \quad Q > 0, \quad q > 0$

Toy model: $A_C = \int [-q\dot{p} - (\frac{1}{2m} p^2 - e^2/q)] dt$

Enhanced classical action: $\hbar > 0 ; \quad \langle \eta | (aQ + bD) | \eta \rangle = a$

$A_R = \int \langle p, q | [i\hbar \partial / \partial t - \mathcal{H}(P, Q)] | p, q \rangle dt = \int [-q\dot{p} - H(p, q)] dt$

$$H(p, q) = \langle p, q | \mathcal{H}(P, Q) | p, q \rangle = \langle \eta | \mathcal{H}(P/q + p, qQ) | \eta \rangle$$

Model: $\langle p, q | \frac{1}{2m} P^2 - e^2/Q | p, q \rangle = \frac{1}{2m} p^2 - C/q + C'/q^2 \quad \leftarrow$

$$C = \langle \eta | e^2/Q | \eta \rangle \propto e^2 ; \quad C' = \frac{1}{2m} \langle \eta | P^2 | \eta \rangle \propto \frac{1}{2m} \hbar^2$$

$$\underline{C' \cong (\hbar^2/2me^2) C \cong (Bohr\ radius) C}$$

Spin States & Enhan. Quan.

Spin coherent states: $[S_1, S_2] = i\hbar S_3$; $[\hbar > 0]$

$$|\theta, \varphi\rangle = e^{-i\varphi S_3/\hbar} e^{-i\theta S_2/\hbar} |s, s\rangle ; \quad S_3 |s, m\rangle = m\hbar |s, m\rangle$$

$$\begin{aligned} A_R &= \int \langle \theta(t), \varphi(t) | [i\hbar \partial / \partial t - \mathcal{H}(\mathbf{S})] | \theta(t), \varphi(t) \rangle \, dt \\ &= \int [s\hbar \cos(\theta(t)) \dot{\varphi}(t) - H(\theta(t), \varphi(t))] \, dt \\ &= \int [p(t) \dot{q}(t) - H(p(t), q(t))] \, dt \end{aligned}$$

Fubini - Study metric:

$$\begin{aligned} d\sigma^2 &= s\hbar[d\theta^2 + \sin(\theta)^2 d\varphi^2] \\ &= (1 - p^2 / s\hbar)^{-1} dp^2 + (1 - p^2 / s\hbar) \, dq^2 \end{aligned}$$

TOPIC 3

- Rotationally symmetric models [$\vec{p}^2 \equiv \vec{p} \cdot \vec{p}$]

$$H(\vec{p}, \vec{q}) = \frac{1}{2}[\vec{p}^2 + m_0^2 \vec{q}^2] + \lambda_0 (\vec{q}^2)^2 , \quad \vec{p} = \{p_n\}_{n=1}^N$$

- Free quantum models for $N \leq \infty$
- Interacting quantum models for $N < \infty$

$$H(\vec{p}, \vec{q}) \equiv \langle \vec{p}, \vec{q} | \mathcal{H} | \vec{p}, \vec{q} \rangle$$

- **Reducible operator representation is the key**

$$H(\vec{p}, \vec{q}) \equiv \langle \vec{p}, \vec{q} | \mathcal{H}(\vec{P}, \vec{Q}, \dots) | \vec{p}, \vec{q} \rangle$$

Rotationally Sym. Models (1)

Phase space coordinates : $\vec{p} = (p_1, \dots, p_N)$, $\vec{q} = (q_1, \dots, q_N)$

$$H(\vec{p}, \vec{q}) = \frac{1}{2}[\vec{p}^2 + m_0^2 \vec{q}^2] + \lambda_0 (\vec{q}^2)^2 ; \quad N \leq \infty$$

Invariant under $\vec{p} \rightarrow O\vec{p}$, $\vec{q} \rightarrow O\vec{q}$; $O \in \text{SO}(N, \mathbb{R})$

Basic invariants: $X \equiv \vec{p}^2$, $Y \equiv \vec{p} \cdot \vec{q}$, $Z \equiv \vec{q}^2$

Constants of motion: E , $\vec{L}^2 = (\vec{p} \times \vec{q})^2 = XZ - Y^2$



Quantization: $\vec{p} \rightarrow \vec{P}$, $\vec{q} \rightarrow \vec{Q}$; $[Q_j, P_k] = i\hbar \delta_{jk}$

Hamiltonian : $\mathcal{H} = \frac{1}{2} : \vec{P}^2 + m_0^2 \vec{Q}^2 : + \lambda_0 : (\vec{Q}^2)^2 :$, $N < \infty$

When $N \rightarrow \infty$, it is necessary that $\lambda_0 = \lambda / N$

Rotationally Sym. Models (2)

Schroedinger equation with a real unique ground state

$\varphi_N(\vec{x})$ with full rotational symmetry: $\psi_N(r) = \varphi_N(\vec{x})$.


Fourier transformation of the ground state distribution

$$\begin{aligned} C_N(\vec{p}) &\equiv \int e^{i\vec{p}\cdot\vec{x}/\hbar} \varphi_N(\vec{x})^2 d\vec{x} \\ &= \int e^{ipr\cos(\theta)/\hbar} \psi_N(r)^2 r^{N-1} \sin(\theta)^{N-2} dr d\theta d\Omega_{N-2} \\ &\cong K_N \int e^{-p^2 r^2 / 2\hbar^2(N-2)} \psi_N(r)^2 r^{N-1} dr d\Omega_{N-2} \\ &\rightarrow \int_0^\infty e^{-bp^2/2\hbar} f(b) db ; \quad [\int_0^\infty f(b) db = 1] \quad \text{sym} \end{aligned}$$

Uniqueness: $f(b) = \delta(b - 1/2m)$

A free theory!

Result: $C_\infty(\vec{p}) = e^{-p^2/4m\hbar}$;



~~L&L~~

Magic Dots

$$:\! e^{i(\alpha P + \beta Q)/\hbar} :\! \equiv e^{i(\alpha P + \beta Q)/\hbar} / \langle \eta | e^{i(\alpha P + \beta Q)/\hbar} | \eta \rangle$$

$$\langle \eta | :\! e^{i(\alpha P + \beta Q)/\hbar} :\! | \eta \rangle = 1 \quad ; \quad | p, q \rangle \equiv e^{i(pQ - qP)/\hbar} | \eta \rangle$$

$$\langle p, q | :\! e^{i(\alpha P + \beta Q)/\hbar} :\! | p, q \rangle \equiv \langle \eta | :\! e^{i(\alpha(P+p) + \beta(Q+q))/\hbar} :\! | \eta \rangle$$

$$\equiv e^{i(\alpha p + \beta q)/\hbar}$$

$$\langle p, q | :\! H(P, Q) :\! | p, q \rangle \equiv H(p, q) \text{ form } \underline{\text{OR}} \text{ operator?}$$

$$\text{operator needs: } \langle p, q | \{ :\! H(P, Q) :\! \}^2 | p, q \rangle < \infty$$

Rotationally Sym. Models (3)

$$(m_0 \vec{Q} + i\vec{P}) |0\rangle = 0 ; \quad |\vec{p}, \vec{q}\rangle = \exp[i(\vec{p} \cdot \vec{Q} - \vec{q} \cdot \vec{P})/\hbar] |0\rangle \quad \text{G}$$

$$\begin{aligned} \langle \vec{p}, \vec{q} | \tilde{\mathcal{H}} | \vec{p}, \vec{q} \rangle &= \langle \vec{p}, \vec{q} | \{ \frac{1}{2} : \vec{P}^2 + m_0^2 \vec{Q}^2 : + w : (\vec{P}^2 + m_0^2 \vec{Q}^2)^2 : \} | \vec{p}, \vec{q} \rangle \\ &= \frac{1}{2} (\vec{p}^2 + m_0^2 \vec{q}^2) + w (\vec{p}^2 + m_0^2 \vec{q}^2)^2 ; \quad N \leq \infty \end{aligned}$$

$$[m(\vec{Q} + \zeta \vec{S}) + i\vec{P}] |0; \zeta\rangle = [m(\vec{S} + \zeta \vec{Q}) + i\vec{R}] |0; \zeta\rangle = 0 ; \quad 0 < \zeta < 1 \quad \text{G}$$

$$|\vec{p}, \vec{q}; \zeta\rangle \equiv \exp[i(\vec{p} \cdot \vec{Q} - \vec{q} \cdot \vec{P})/\hbar] |0; \zeta\rangle$$

$$\begin{aligned} \langle \vec{p}, \vec{q}; \zeta | \tilde{\mathcal{H}} | \vec{p}, \vec{q}; \zeta \rangle &= \langle \vec{p}, \vec{q}; \zeta | \{ \frac{1}{2} : \vec{P}^2 + m^2 (\vec{Q} + \zeta \vec{S})^2 : \\ &\quad + \frac{1}{2} : \vec{R}^2 + m^2 (\vec{S} + \zeta \vec{Q})^2 : + v : [\vec{R}^2 + m^2 (\vec{S} + \zeta \vec{Q})^2]^2 : \} | \vec{p}, \vec{q}; \zeta \rangle \\ &= \frac{1}{2} (\vec{p}^2 + m^2 \vec{q}^2) + \frac{1}{2} \zeta^2 m^2 \vec{q}^2 + v \zeta^4 m^4 (\vec{q}^2)^2 \\ &\equiv \frac{1}{2} (\vec{p}^2 + m_0^2 \vec{q}^2) + \lambda_0 (\vec{q}^2)^2 ; \quad N \leq \infty \end{aligned}$$

T
E
S
T

R
E
A
L



Rot. Sym. Models : Summary

- Conventional quantization works if N is finite but leads to ***triviality*** if N is infinite
- *Enhanced quantization applies even for reducible operator representations*
- Using the ***Weak Correspondence Principle*** $H(\vec{p}, \vec{q}) = \langle \vec{p}, \vec{q} | \mathcal{H} | \vec{p}, \vec{q} \rangle$
a *nontrivial* quantization results if N is finite or N is infinite --- with ***NO divergences!***
- *Class. & Quant. formalism is similar for all N*



TOPIC 4

- A 3 + 1 Einstein gravity formalism

$$A = \int \{ \pi^{ab}(t, x) \dot{g}_{ab}(t, x) - N^a(t, x) H_a(t, x) - N(t, x) H(t, x) \} dt d^3x$$

- Metric positivity $u^a g_{ab}(t, x) u^b > 0$
- Problems with canonical variables
- Relevance of affine variables
- Imposition of constraints
- Functional integral formulation

Choice of Classical Variables

Classical variables: $x \in \mathbf{R}^3$

$$\pi_c^a(x) = \pi^{ab}(x)g_{bc}(x) ; \quad g_{ab}(x) , \quad \underline{u^a g_{ab}(x) u^b > 0}$$

$$\{g_{ab}(x), \pi^{cd}(y)\} = \frac{1}{2}[\delta_a^c \delta_b^d + \delta_a^d \delta_b^c] \delta(x, y)$$

and

$$\{\pi_b^a(x), \pi_d^c(y)\} = \frac{1}{2}[\delta_d^a \pi_b^c(x) - \delta_b^c \pi_d^a(x)] \delta(x, y)$$

$$\{g_{ab}(x), \pi_d^c(y)\} = \frac{1}{2}[\delta_a^c g_{bd}(x) + \delta_b^c g_{ad}(x)] \delta(x, y)$$

$$\{g_{ab}(x), g_{cd}(y)\} = 0$$

Choice of Quantum Variables

$$[\hat{g}_{ab}(x), \hat{\pi}^{cd}(y)] = \frac{i}{2} [\delta_a^c \delta_b^d + \delta_a^d \delta_b^c] \delta(x, y)$$

or

$$[\hat{\pi}_b^a(x), \hat{\pi}_d^c(y)] = \frac{i}{2} [\delta_d^a \hat{\pi}_b^c(x) - \delta_b^c \hat{\pi}_d^a(x)] \delta(x, y)$$

$$[\hat{g}_{ab}(x), \hat{\pi}_d^c(y)] = \frac{i}{2} [\delta_a^c \hat{g}_{bd}(x) + \delta_b^c \hat{g}_{ad}(x)] \delta(x, y)$$

c.c.r.

$$[\hat{g}_{ab}(x), \hat{g}_{cd}(y)] = 0$$

$$U(\gamma) = \exp[-i \int \gamma_a^b \hat{\pi}_b^a d^3 y] , \quad M_a^c(x) = \{\exp[\gamma(x)/2]\}_a^c$$

$$U(\gamma)^{-1} \hat{g}_{ab}(x) U(\gamma) = M_a^c(x) \hat{g}_{cd}(x) M_b^d(x)$$

Coh. States of Quantum Gravity

$$|\pi, g\rangle = e^{i\int \pi^{cd} \hat{g}_{cd} d^3y} e^{-i\int \gamma_a^b \hat{\pi}_b^a d^3y} |\eta\rangle$$

$$\langle \eta | \hat{g}_{cd}(x) | \eta \rangle = \tilde{g}_{cd}(x) , \quad \langle \eta | \hat{\pi}_b^a(x) | \eta \rangle = 0$$

$$\begin{aligned} \langle \pi'', g'' | \pi', g' \rangle &= \exp \left\{ -2 \int b(x) d^3x \right. \\ &\times \left. \ln \left(\frac{\det\{(g''^{ab} + g'^{ab})/2 + i b(x)^{-1} (\pi''^{ab} - \pi'^{ab})/2\}}{[\det\{g''^{ab}\} \det\{g'^{ab}\}]^{1/2}} \right) \right\} \end{aligned}$$

$$g_{ab}(x) = M_a^c(x) \tilde{g}_{cd}(x) M_b^d(x) , \quad M_a^c(x) = \{\exp[\gamma(x)/2]\}_a^c$$

Formal Path Integrals

Conventional quantization :

$$Q|q\rangle = q|q\rangle , \quad P|p\rangle = p|p\rangle \quad ; \quad (\hbar = 1)$$

$$\langle q''|q'\rangle = \delta(q''-q') = \int e^{ip(q''-q')} dp / 2\pi$$

$$= \mathcal{M} \int e^{i \int p(t) \dot{q}(t) dt} DpDq$$



$$\langle q'', t'' | q', t' \rangle = \langle q'' | e^{-i \mathcal{H}(t''-t')} | q' \rangle$$

$$= \mathcal{M} \int e^{i \int [p(t) \dot{q}(t) - H(p(t), q(t))] dt} DpDq$$



Genuine Path Integrals-1

Canonical coherent state case: $| p, q \rangle = e^{-iqP} e^{ipQ} | 0 \rangle$

$$\langle p'', q'' | p', q' \rangle = e^{i(p''+p')(q''-q')'2 - [(p''-p')^2 + (q''-q')^2]/4}$$

$$= \lim_{\nu \rightarrow \infty} \mathcal{N} \int e^{i \int p(t) \dot{q}(t) dt} e^{- \int [\dot{p}(t)^2 + \dot{q}(t)^2] dt / 2\nu} Dp Dq$$

$$= \lim_{\nu \rightarrow \infty} 2\pi e^{\nu(t''-t')/2} \int e^{i \int p(t) \dot{q}(t) dt} d\mu_\nu^C(p, q)$$



$$\langle p'', q'', t'' | p', q', t' \rangle = \langle p'', q'' | e^{-i\mathcal{H}(t''-t')} | p', q' \rangle$$

$$= \lim_{\nu \rightarrow \infty} \mathcal{N} \int e^{i \int p(t) \dot{q}(t) dt - h(p(t), q(t))] dt}$$

$$\times e^{- \int [\dot{p}(t)^2 + \dot{q}(t)^2] dt / 2\nu} Dp Dq$$



$$= \lim_{\nu \rightarrow \infty} 2\pi e^{\nu(t''-t')/2} \int e^{i \int p(t) \dot{q}(t) dt - h(p(t), q(t))] dt} d\mu_\nu^C(p, q)$$

Genuine Path Integrals-2

Affine coherent state case : $|p, q\rangle = e^{ipQ} e^{-i\ln(q)D} |\beta\rangle$, $q > 0$

$$\begin{aligned}\langle p'', q'' | p', q' \rangle &= \{\tfrac{1}{2}[(q''/q')^{\frac{1}{2}} + (q'/q'')^{\frac{1}{2}} + (i/\beta)(q''q')^{\frac{1}{2}}(p''-p')] \}^{-2\beta} \\ &= \lim_{\nu \rightarrow \infty} \mathcal{N} \int e^{-i \int q(t) \dot{p}(t) dt} e^{-\int [\beta^{-1}q(t)^2 \dot{p}(t)^2 + \beta q(t)^{-2} \dot{q}(t)^2] dt / 2\nu} Dp Dq \\ &= \lim_{\nu \rightarrow \infty} 2\pi e^{\nu(t''-t')/2} \int e^{-i \int q(t) \dot{p}(t) dt} d\mu_\nu^A(p, q)\end{aligned}$$



$$\langle p'', q'', t'' | p', q', t' \rangle = \langle p'', q'' | e^{-i\mathcal{H}(t''-t')} | p', q' \rangle$$

$$\begin{aligned}&= \lim_{\nu \rightarrow \infty} \mathcal{N} \int e^{i \int [-q(t) \dot{p}(t) dt - h(p(t), q(t))] dt} \\ &\quad \times e^{-\int [\beta^{-1}q(t)^2 \dot{p}(t)^2 + \beta q(t)^{-2} \dot{q}(t)^2] dt / 2\nu} Dp Dq\end{aligned}$$



$$= \lim_{\nu \rightarrow \infty} 2\pi e^{\nu(t''-t')/2} \int e^{i \int [-q(t) \dot{p}(t) dt - h(p(t), q(t))] dt} d\mu_\nu^A(p, q)$$

Quantum Constraints

Constraints: $\varphi_\alpha(p, q) = 0$, $\sum_{\alpha=1}^A \varphi_\alpha(p, q)^2 = 0$


Dirac: $\Phi_\alpha(P, Q) |\psi_{\text{phys}}\rangle = 0$, (limited use)

POM: $\mathbf{E} = \mathbf{E}(\sum_\alpha \Phi_\alpha^2 \leq \delta(\hbar)^2)$, $\mathbf{H}_{\text{phys}} = \mathbf{E}\mathbf{H}$

$$(1) \quad \mathbf{E}(J_1^2 + J_2^2 + J_3^2 \leq \hbar^2 / 2)$$

$$(2) \quad \mathbf{E}(P^2 + Q^2 \leq \hbar)$$

$$(3) \quad \mathbf{E}(Q^2 \leq \delta^2) = \mathbf{E}(-\delta < Q < \delta)$$



$$\lim_{\delta \rightarrow 0} \langle p'', q'' | \mathbf{E} | p', q' \rangle / \langle \eta | \mathbf{E} | \eta \rangle = \langle \langle p'', q'' | p', q' \rangle \rangle$$

→ $\mathbf{E}(\sum_\alpha \Phi_\alpha^2 \leq \delta(\hbar)^2) = \int \mathbf{T} e^{-i \int \lambda^\alpha(t) \Phi_\alpha dt} DR(\lambda)$ ← 59

Functional Integral Reps.

$$\langle \pi'', g'' | \pi', g' \rangle$$

$$\begin{aligned}
&= \lim_{\nu \rightarrow \infty} M_\nu \int \exp \{-i \int [g_{ab} \dot{\pi}^{ab}] dt d^3x\} \\
&\times \exp \{-(1/2\nu) \int [b^{-1}(x) g_{ab} g_{cd} \dot{\pi}^{bc} \dot{\pi}^{da} + b(x) g^{ab} g^{cd} \dot{g}_{bc} \dot{g}_{da}] dt d^3x\} \\
&\times [\prod_{t,x} \prod_{a \geq b} d\pi^{ab}(t,x) dg_{ab}(t,x)]
\end{aligned}$$

$$\langle \pi'', g'' | \mathbf{E} | \pi', g' \rangle$$

$$\begin{aligned}
&= \lim_{\nu \rightarrow \infty} M_\nu \int \exp \{-i \int [g_{ab} \dot{\pi}^{ab} + N^a H_a + NH] dt d^3x\} \\
&\times \exp \{-(1/2\nu) \int [b^{-1}(x) g_{ab} g_{cd} \dot{\pi}^{bc} \dot{\pi}^{da} + b(x) g^{ab} g^{cd} \dot{g}_{bc} \dot{g}_{da}] dt d^3x\} \\
&\times [\prod_{t,x} \prod_{a \geq b} d\pi^{ab}(t,x) dg_{ab}(t,x)] DR(N^a, N)
\end{aligned}$$

Affine Quan. Gravity : Summary

- Metric positivity is preserved by use of affine variables
- Constraints may be imposed by the projection operator method
- Phase-space appearing functional integrals lead to affine quantization
- Affine approach has potential to eliminate conventional divergences

TOPIC 5

- Ashtekar, Henderson, and Sloan model
- Singularity, omitting spatial derivatives (BKL)
- Relevance of affine variables
- Imposition of Hamiltonian constraint
- Classical solutions exhibit “spikes”
- Do quantum solutions have “spikes”, or not

Quantum Spikes? (1)

Ashtekar, Henderson, and Sloan model

variables: $(C_I, P^J) = (C_I, -2D^J)$, $(I, J) \in (1, 2, 3)$; φ, π

$$C := C_1 + C_2 + C_3, \quad D := D^1 + D^2 + D^3$$

Poisson brackets (affine Lie algebra):

$$\{C_I, C_J\} = 0 = \{D^I, D^J\}, \quad \{C_I, D^J\} = \delta_I^J C_J ; \quad \{\varphi, \pi\} = 1$$

classical Hamiltonian : $H = \frac{1}{2}C^2 - \sum_I C_I^2 + 2D^2 - 4\sum_I D^{I2} - \frac{1}{2}\pi^2$

classical constraint: $H = 0$

affine quantization: $\psi \in L^2(\mathbf{R}^4, d^3x \, d\varphi)$, $\pi \rightarrow -i\hbar\partial/\partial\varphi$

$$C_I \rightarrow x_I, \quad D^I \rightarrow -(i\hbar/2)[x_I(\partial/\partial x_I) + (\partial/\partial x_I)x_I]$$

A natural case for an affine quantization !



Quantum Spikes? (2)

Ashtekar, Henderson, and Sloan model

affine quantization : $\psi \in L^2(\mathbf{R}^4, d^3x \, d\varphi) =: \mathbf{H}$

quantum Hamiltonian :

$$\begin{aligned}\mathcal{H} = & \frac{1}{2} \{ \sum_I x_I \}^2 - \sum_I x_I^2 - \frac{1}{2} \hbar^2 \{ \sum_I [x_I (\partial / \partial x_I) + (\partial / \partial x_I) x_I] \}^2 \\ & + \hbar^2 \{ \sum_I [x_I (\partial / \partial x_I) + (\partial / \partial x_I) x_I]^2 \} + \frac{1}{2} \hbar^2 \partial^2 / \partial \varphi^2\end{aligned}$$

quantum constraint: $\mathcal{H}\Psi_{\text{phys}} = 0$

physical Hilbert space: $\mathbf{H}_{\text{phys}} := \mathbf{E}\mathbf{H}$

$$(\mathbf{E} = \mathbf{E}^* = \mathbf{E}^2) , \quad \mathbf{E} = \mathbf{E}(\mathcal{H} = 0)$$

A natural case for an affine quantization !



Quantum Spikes? (3)

Ashtekar, Henderson, and Sloan model ($\varphi = 0 = \pi$) , $[\hbar = 1]$

$$p = (p^1, p^2, p^3) , \quad q = (q_1, q_2, q_3) ; \quad (q > 0)$$

affine coherent states: $|p, q\rangle = e^{ip \cdot \mathbf{C}} e^{-i \ln(q) \cdot \mathbf{D}} |\beta\rangle$, $\langle \beta | (a\mathbf{C} + b\mathbf{D}) | \beta \rangle = a$

$$[\mathbf{C} - 1 + (i/\beta)\mathbf{D}] |\beta\rangle = 0 \Rightarrow \langle p, q | \mathbf{C} | p, q \rangle = q, \langle p, q | \mathbf{D} | p, q \rangle = d$$

enhanced classical Hamiltonian :

$$d = (d^1, d^2, d^3) := (p^1 q_1, p^2 q_2, p^3 q_3)$$

$$q \mathbf{C} := (q_1 \mathbf{C}_1, q_2 \mathbf{C}_2, q_3 \mathbf{C}_3) , \quad d \mathbf{C} := (d^1 \mathbf{C}_1, d^2 \mathbf{C}_2, d^3 \mathbf{C}_3)$$

$$\begin{aligned} H(q, d) &= \langle p, q | \mathcal{H}(\mathbf{C}, \mathbf{D}) | p, q \rangle = \langle \beta | \mathcal{H}(q \mathbf{C}, \mathbf{D} + d \mathbf{C}) | \beta \rangle \\ &= \mathcal{H}(q, d) + \mathcal{O}(\hbar; p, q) \end{aligned}$$

A natural case for an affine quantization !



Quantum Spikes? (4)

Ashtekar, Henderson, and Sloan model ($\pi = \varphi = 0$)

affine quantization: $\psi \in L^2(\mathbf{R}^3, d^3x) = \mathbf{H}$, $\Psi = \Psi_{\text{phys}} \in \mathbf{H}_{\text{phys}}$

quantum Hamiltonian: $\mathcal{H} = \mathcal{H} - \frac{3}{2}\hbar^2$

$\mathcal{H} = \frac{1}{2}[\sum_{I \neq J} x_I x_J - \sum_I x_I^2] - 2\hbar^2 [\sum_{I \neq J} x_I x_J (\partial^2 / \partial x_I \partial x_J) - \sum_I x_I^2 (\partial^2 / \partial x_I^2)]$

quantum constraint: $\mathcal{H}\Psi(x) = 0$; $\theta(y) = 1, y > 0 = 0, y < 0$

$$|J_{+,I}|^2 + |J_{-,I}|^2 = 1 = |K_{+,I}|^2 + |K_{-,I}|^2$$

$$\Psi(x) = \hbar^{-3/2} \prod_I [J_{+,I} \theta(x_I) + J_{-,I} \theta(-x_I)] \exp[-|x_I|/2\hbar]$$

$$\Phi(x) = \hbar^{-3/2} \prod_I [K_{+,I} \theta(x_I) + K_{-,I} \theta(-x_I)] \exp[-|x_I|/2\hbar]$$

EIGHT INDEPENDENT SOLUTIONS

A natural case for an affine quantization !



Quantum Spikes? (5)

Ashtekar, Henderson, and Sloan model ($\pi = \varphi = 0$) , $C_1 = C_1(0)$

$$C_1(t) = e^{i\mathcal{H}t/\hbar} C_1 e^{-i\mathcal{H}t/\hbar} = C_1 + (it/\hbar)[\mathcal{H}, C_1] + \frac{1}{2}(it/\hbar)^2[\mathcal{H}, [\mathcal{H}, C_1]] + \dots$$

$(\Phi, C_1(t)\Psi) = (\Phi, M(t, C_I)\Psi)$, $M(t, C_I)$ is an 8×8 matrix

position in space $\bar{x}_1 = (\Psi[\bar{x}_1], C_1\Psi[\bar{x}_1])$; $\bar{x}_1 = 0$, $\bar{x}_1 > 0$

$$\Psi[0] = 2^{-1/2} \hbar^{-1/2} [\theta(x_1) + \theta(-x_1)] \exp[-|x_1|/2\hbar]$$

$$\begin{aligned} \Psi[\bar{x}_1] = 2^{-1/2} \hbar^{-1/2} & [(1 + \bar{x}_1/\hbar)^{1/2} \theta(x_1) + (1 - \bar{x}_1/\hbar)^{1/2} \theta(-x_1)] \\ & \times \exp[-|x_1|/2\hbar] \end{aligned}$$

$$|(\Psi[\bar{x}_1], C_1(t)\Psi[\bar{x}_1]) - (\Psi[0], C_1(t)\Psi[0])| \leq 2 \|\Psi[\bar{x}_1] - \Psi[0]\| \|M(t, C_I)\|$$

THEREFORE NO QUANTUM SPIKES


A natural case for an affine quantization !



Over All Summary



- Enhanced quantization leads to $C \subset Q$
- Affine variables may be used instead of canonical variables in EQ
- Affine and canonical quantization can lead to different results in EQ
- Affine analysis of ultralocal scalar models succeeds, while canonical analysis fails
- Affine methods are natural for gravity
- Affine methods say NO quantum “spikes”⁶⁸

Other Enh. Quant. Projects

- Additional sheets of vectors in Hilbert space relating quan. and class. models (done) 
- Covariant scalar models φ_n^4 (done) 
- Nonrenormalizable scalar fields (done) 
- Simple models of affine quantization eliminating classical singularities (on going)
- Incorporating constrained systems within enhanced quantization (partially)
- Affine quantum gravity (partially)
- Extension to fermion fields (hints)

Main Message of Today

**Class./Quan. Coexistence *IS*
Possible *AND IT IS* Beneficial**



GOOD

BETTER





References

- ``Enhanced Quantization: A Primer'', *J. Phys. A: Math. Theor.* **45**, 285304 (8pp) (2012);
[arXiv:1204.2870](https://arxiv.org/abs/1204.2870)
- ``Enhanced Quantum Procedures that Resolve Difficult Problems''; *Rev. Math. Physics* 27, 1530002 (43pp) (2015); [arXiv: 1206.4017](https://arxiv.org/abs/1206.4017)
- Ultralocal Model Scalar Quantum Fields:
“*Beyond Conventional Quantization*”
(Cambridge, 2000 & 2005)
- Covariant Scalar Models and Affine Quantum Gravity, “*Enhanced Quantization: Particles, Fields & Gravity*” (World Scientific, 2015)



References - 1

- ``Enhanced Quantization: A Primer'', *J. Phys. A: Math. Theor.* **45**, 285304 (8pp) (2012);
[arXiv:1204.2870](#)
- ``Enhanced Quantum Procedures that Resolve Difficult Problems''; *Rev. Math. Physics* 27, 1530002 (43pp) (2015); [arXiv: 1206.4017](#)
- ``Revisiting Canonical Quantization''; *Mod. Phys. Lett. A*, 29, 1430020; [arXiv: 1211.735](#)
- “Affine Quantization and the Initial Cosmological Singularity”, M. Fanuel and S. Zonetti, *EPL*, 101, 10001 (2013); [arXiv:1203.4936v3](#)

References - 2

- ``Scalar Field Quantization Without Divergences In All Spacetime Dimensions'' *J. Phys. A: Math. Theor.* **44**, 273001 (2011); [arXiv:1101.1706](#)
- ``Divergences in Scalar Quantum Field Theory: The Cause and the Cure'', *Mod. Phys. Lett. A* **27**, 1250117 (9pp) (2012); [arXiv:1112.0803](#)
- Ultralocal Model Scalar Quantum Fields:
“Beyond Conventional Quantization”
(Cambridge, 2000 & 2005)
- Covariant Scalar Models and Affine Quantum Gravity, *“Enhanced Quantization: Particles, Fields & Gravity”* (World Scientific, 2015)



Other Ultralocal Stories

Early attempts at an ultralocal model:

classical model as a classical mechanical model

$$H = \int \{ \frac{1}{2} [\pi(x)^2 + m^2 \varphi(x)^2] + g \varphi(x)^4 \} d^s x$$

$$H_N = \sum_k \{ \frac{1}{2} [\pi_k^2 + m^2 \varphi_k^2] + g \varphi_k^4 \} a^s \rightarrow H$$



$$\hat{H}_N = -\frac{1}{2} a^{-s} \hbar^2 \sum_k \frac{\partial^2}{\partial \phi_k^2} + \frac{1}{2} m_0^2 \sum_k \phi_k^2 a^s + g_0 \sum_k \phi_k^4 a^s - E_0$$

strong coupling gravity (M. Palati)

$$H_{scg} = \int \{ g(x)^{-1/2} [\pi_b^a(x) \pi_a^b(x) - \frac{1}{2} \pi_a^a(x) \pi_b^b(x)] + 2 \Lambda_C g(x)^{1/2} \} d^s x$$




Does $[Q, R] = 0$? (No!)

$$I = \int [p\dot{q} - \lambda(p^2 + q^4 - E)]dt$$

What values of (positive) E are allowed?

$$\rightarrow [(P^2 + Q^4) - E] |\psi\rangle = 0 , \quad \{E_n\}$$

$$\begin{aligned}\rightarrow & \int \delta\{p^2 + q^4 - E\} e^{i\int p\dot{q}dt} DpDq \\ & \int \delta\{p\} \Pi(4q^3) \delta\{p^2 + q^4 - E\} e^{i\int p\dot{q}dt} DpDq \\ & = \int \Pi(4q^3) \delta\{q^4 - E\} Dq \\ & = 1 \text{ or } 0 , \quad \text{independent of } E !\end{aligned}$$



Positive or Non-negative ?

$$\int_{q>0} e^{i\int [p\dot{q}-qp^2]dt} DpDq \\ = \int_{q>0} e^{i\int [\dot{q}^2/4q]dt} \Pi_t dq / \sqrt{2q}$$

Let $u^2 = 2q$, $du = dq / \sqrt{2q}$, $\dot{u}^2 = \dot{q}^2 / 2q$

$$= \int_{u\neq 0} e^{i\int \dot{u}^2 / 2 dt} Du \quad (\text{forget } u=0)$$

$$= \int e^{i\int \dot{u}^2 / 2 dt} Du = \text{free particle!} \quad \text{Wrong answer!}$$

$$= \int e^{i\int (\dot{u}_1^2 + \dot{u}_2^2) / 2 dt} Du_1 Du_2 d\theta_{12} / 2\pi$$

Right answer!